

For control of piercing and sucking insects in commercial production of greenhouse-grown cucumbers, fruiting vegetables, strawberries, and strawberry transplants grown for the home consumer market

This supplemental label expires October 30, 2023, and must not be used or distributed after this date.

#### **Active Ingredient:**

Afidopyropen, [(3S,4R,4aR,6S,6aS,12R,12aS,12bS)-3-[(cyclopropylcarbonyl)oxy]-1,3,4,4a,5,6,6a,12,12a,12b-decahydro-6,12-dihydroxy-4,6a,12b-trimethyl-11-oxo-9-(3-pyridinyl)-2H,11H-naphtho[2,1-b]pyrano[3,4-e]pyran-4-yl]methyl cyclopropanecarboxylate

 cyclopropanecarboxylate
 9.78%

 Other Ingredients:
 90.22%

 Total:
 100.00%

Contains 0.83 pound of afidopyropen per gallon, formulated as a dispersible concentrate.

# EPA Reg. No. 7969-393 CAUTION/PRECAUCION

#### **Directions For Use**

- It is a violation of federal law to use this product in a manner inconsistent with its labeling.
- The supplemental labeling and the entire
   Ventigra® insecticide container label, EPA Reg. No. 7969-393, must be in possession of the user at the time of application.
- Read the label affixed to the container for Ventigra before applying.

 Use of Ventigra according to this labeling is subject to the use precautions and limitations imposed by the label affixed to the container for Ventigra.

# **Application Instructions**

Refer to the **Ventigra** container label for additional application instructions and restrictions.



### **Use Sites**

**Ventigra** can be used on the following greenhouse crops grown to production (edible fruit):

- Berry, low growing
  - Strawberry
- Cucurbit Vegetables
  - Cucumber
- Vegetable, Fruiting, Group 8-10
  - African eggplant
  - Bush tomato
  - Cocona
  - Currant tomato
  - Eggplant
  - Garden huckleberry
  - Goji berry
  - Groundcherry
  - Martynia
  - Naranjilla
  - Okra
  - Pea eggplant
  - Pepino
  - Pepper, bell
  - Pepper, nonbell
  - Roselle
  - Scarlet eggplant
  - Sunberry
  - Tomatillo
  - Tomato
  - Tree tomato
  - Cultivars, varieties, and/or hybrids of these commodities

**Ventigra® insecticide** may also be used on the following greenhouse-grown transplants for the home consumer market only:

- Berry, low growing
  - Strawberry

Table 1. Ventigra® insecticide Use-specific Application Instructions

| Сгор                                                                                                                                                                                                                       | Pest                                                                                                                                   | Product Use<br>Rate per<br>Application<br>(fl ozs/100<br>gallons) | Minimum<br>Time from<br>Application<br>to Harvest<br>(PHI) days | Use Directions                                                                                                                                            |
|----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|----------------------------------------------------------------------------------------------------------------------------------------|-------------------------------------------------------------------|-----------------------------------------------------------------|-----------------------------------------------------------------------------------------------------------------------------------------------------------|
| Berry, Low Growing<br>Strawberry                                                                                                                                                                                           | Aphids such as:<br>Cotton/Melon aphid<br>(Aphis spp.)                                                                                  | 1.4 (aphids)<br>7.0 (psyllids<br>and whiteflies)                  | 0                                                               | Ventigra is not a rescue treatment and should be applied at the onset of pest infestation.                                                                |
| Cucurbit Vegetables Cucumber Cultivars, varieties, and/or hybrids of these Vegetable, Fruiting,                                                                                                                            | Green peach aphid (Myzus spp.) Rose aphid (Macrosiphum spp.) Strawberry aphid (Chaetosiphon spp.)                                      |                                                                   |                                                                 | For use in commercial production of strawberries and for strawberry transplants for the home consumer market.                                             |
| Group 8-10 African eggplant                                                                                                                                                                                                |                                                                                                                                        |                                                                   |                                                                 | <b>DO NOT</b> make applications at intervals shorter than 7 days.                                                                                         |
| Bush tomato Cocona Currant tomato Eggplant Garden huckleberry                                                                                                                                                              | Psyllids such as: Potato/tomato psyllid (Bactericera cockerelli) Whiteflies such as:                                                   |                                                                   |                                                                 | DO NOT apply more than the maximum seasonal use rate of 14 fl ozs of <b>Ventigra</b> (0.09 lb afidopyropen ai) per acre per crop season.                  |
| Goji berry Groundcherry Martynia Naranjilla Okra Pea eggplant Pepino Pepper, bell Pepper, nonbell Roselle Scarlet eggplant Sunberry Tomatillo Tomato Tree tomato Cultivars, varieties, and/or hybrids of these commodities | Greenhouse whitefly ( <i>Trialeurodes</i> spp.) Silverleaf whitefly ( <i>Bemisia</i> spp.) Sweetpotato whitefly ( <i>Bemisia</i> spp.) |                                                                   |                                                                 | DO NOT apply more than the annual maximum use rate of 42 fl ozs of Ventigra (0.27 lb afidopyropen ai) per acre per calendar year.                         |
|                                                                                                                                                                                                                            |                                                                                                                                        |                                                                   |                                                                 | Resistance Management. DO NOT make more than 2 sequential applications of Ventigra before using an effective insecticide with a different mode of action. |
|                                                                                                                                                                                                                            |                                                                                                                                        |                                                                   |                                                                 | See the container label for <b>Table</b> 3. Dilution Rate Conversion for Ventigra for information on use rates and respective spray volumes.              |
|                                                                                                                                                                                                                            |                                                                                                                                        |                                                                   |                                                                 | When an adjuvant is to be used with this product, BASF recommends the use of a Chemical Producers and Distributors Association certified adjuvant.        |

<sup>&</sup>lt;sup>1</sup> Silverleaf or Sweetpotato whitefly - *Bemisia tabaci*. *Bemisia tabaci* includes both B and Q biotype.

### **Conditions of Sale and Warranty**

The **Directions For Use** of this product reflect the opinion of experts based on field use and tests. The directions are believed to be reliable and must be followed carefully. However, it is impossible to eliminate all risks inherently associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or use of the product in a manner inconsistent with its labeling, all of which are beyond the control of BASF CORPORATION ("BASF") or the Seller. To the extent consistent with applicable law, all such risks shall be assumed by the Buyer.

BASF warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes referred to in the **Directions For Use**, subject to the inherent risks, referred to above.

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BASF MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF FITNESS OR MERCHANTABILITY OR ANY OTHER EXPRESS OR IMPLIED WARRANTY.

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BUYER'S EXCLUSIVE REMEDY AND BASF'S EXCLUSIVE LIABILITY, WHETHER IN CONTRACT, TORT, NEGLIGENCE, STRICT LIABILITY, OR OTHERWISE, SHALL BE LIMITED TO REPAYMENT OF THE PURCHASE PRICE OF THE PRODUCT.

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BASF AND THE SELLER DISCLAIM ANY LIABILITY FOR CONSEQUENTIAL, EXEMPLARY, SPECIAL OR INDIRECT DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT.

BASF and the Seller offer this product, and the Buyer and User accept it, subject to the foregoing **Conditions of Sale and Warranty** which may be varied only by agreement in writing signed by a duly authorized representative of BASF.

Ventigra is a registered trademark of BASF.

© 2020 BASF Corporation All rights reserved.

007969-00393.20181221f.NVA 2018-04-533-0119

BASF Corporation 26 Davis Drive Research Triangle Park, NC 27709

